

学校编码: 10384
学号: 21220051403186

分类号 _____ 密级 _____
UDC _____

博 士 学 位 论 文

典型海洋环境中浮游细菌多样性 及环境适应机制的研究

Microbial diversity and Environmental Adaptation Mechanisms in Typical Marine Environments

曾 永 辉

指导教师姓名: 焦念志 教授

专 业 名 称: 海 洋 生 物 学

论文提交日期: 2008 年 08 月

论文答辩时间: 2008 年 08 月

学位授予日期: 2008 年 月

答辩委员会主席: 刘康克 教授

评 阅 人: “双盲” 评审

2008 年 08 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其他方式合理复制学位论文。

本学位论文属于：

() 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

() 2、不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

典型海洋环境中浮游细菌多样性及环境适应机制的研究

摘 要

海洋浮游细菌是海洋生态系统中丰度最高的微型生物类群，具有极为丰富的遗传多样性和生理代谢多样性，从而具有极为重要的生态地位和功能。本论文阐明了一些典型的海洋浮游细菌类群在中国的典型海域及全球主要大洋中的多样性分布特征，并探讨了它们的环境适应机制以及和环境协同进化的关系。主要研究对象为受关注较少同时对海洋碳循环和光利用有独特贡献的几类浮游细菌类群，包括擅长降解大分子颗粒有机物的嗜纤维菌—黄杆菌类群（*Cytophaga-Flavobacteria*, CF）、生理潜能和生态功能目前仍知之甚少的浮游古菌、具有固定 CO₂ 能力的浮游变形细菌、能兼性利用光营混合营养的好氧不产氧光合异养细菌（AAPB）和有色异养细菌（PHB）等。

首先，我们选取了东海的长江口为中国边缘海中河口到陆架的典型过渡区域，系统地调查了 CF 类群、浮游古菌和固碳变形细菌在该区域的多样性分布模式。CF 类群是海洋中最为丰富的浮游细菌类群之一。我们通过使用一对新设计的 CF 特异的 16S rRNA 基因引物构建得到长江口与东海两个海区的克隆文库，揭示了 CF 在这两个区域丰富的多样性以及截然不同的多样性分布模式。70 条 CF 的 16S rRNA 基因序列可分为 26 个亚群，包括 7 个东海特有的亚群、17 个长江口特有的亚群以及 2 个共有的亚群。东海 CF 种的多样性较高，而长江口 CF 亚群的多样性较高。在亚群水平上，长江口和东海都是广谱的亚群占优势地位。来源于土壤、近岸和淡水的亚群仅在长江口有分布，这和该站位的地理位置和环境特征是一致的。CF 种对水文条件完全不同的两种生境的适应以及本土 CF 群落的发展和演替造成了长江口与东海 CF 群落结构间的显著差异。

通过构建长江口古菌 16S rRNA 基因的 2 个克隆文库，我们也揭示出浮游古菌这类在海洋生态系统中广泛分布的类群同样也存在于中国的典型海域。对所得到的 21 个 RFLP 带型测序的结果显示了长江口区域存在着两个古菌类群，即属于 *Crenarchaeota* 的海洋类群 I (MG I) 和属于 *Euryarchaeota* 的海洋类群 II (MG II)。MG I 的克隆子在 2 个文库中都是优势类群。我们得到的大部分序列与未培养的海洋古菌近缘，其中有 2 条序列与新鉴定的海洋硝化古菌 *Nitrosopumilus maritimus*

有 98% 的同源性，提示古菌在河口生态系统中可能有着重要的生态作用。

具有固碳潜力的变形细菌也是一类具有重要生态意义但受关注较少的海洋浮游细菌类群。我们首次设计得到专门针对变形细菌 1, 5-二磷酸核酮糖羧化酶/加氧酶 (RubisCO) 大亚基 (*rbcL*) 基因 I 型和 II 型的两套引物，并用于东海近岸和外海两个站位样品的扩增。结果表明 I 型的多样性在高盐低生产力的离岸站位较高，而 II 型的多样性在低盐高生产力的近岸站位较高。部分序列与数据库中已知的 *rbcL* 基因序列相似性很低 (60~78%)。揭示出在东海水域分布着多样的具有固碳能力的浮游变形细菌，这也提示我们需要进一步关注它们在海洋碳循环中的贡献。

然后，我们以一类典型的功能海洋细菌类群 AAPB 为代表，系统地分析了它们在全球主要大洋中的多样性分布模式，并探讨了它们和环境的协同进化关系。来自包括太平洋、大西洋和印度洋以及中国边缘海的表层 AAPB 指示基因——光反应中心复合体小亚基基因 (*pufM*) 的多样性数据表明，AAPB 在全球海洋中具有丰富的多样性，而且随着叶绿素 *a* 浓度的升高而呈降低的趋势。这一发现说明从寡营养到富营养海区，AAPB 的多样性和丰度的分布趋势正好相反。

在垂直梯度上，通过对中心太平洋、大西洋和印度洋三个站位的表层和弱光层上层的浮游细菌样品建立 6 个 *pufM* 基因克隆文库，我们进一步证实了贫营养大洋 AAPB 种群极高的多样性。同时，首次从真光层以下的弱光区域 (200 m) 扩增得到大量的 *pufM* 序列 (共 136 条序列，划分为 37 个 OTU)。系统发育分析的结果表明真光层以下的 AAPB 种群也是多样化的，覆盖了所有在表层海水中发现的亚群，但多样性、GC 和 GC₃ 含量比表层站位稍低，同时优势度更高。这些结果支持了这个假说——AAPB 能利用弱光层的弱光来获得能量，从而可以分布到真光层以下的水层中。

为进一步探讨不产氧光合细菌与环境协同进化的关系，我们从公共数据库收集了 89 个含有 *pufM* 序列的不产氧光合细菌及其来源环境相关的信息。系统发育分析表明 21 个 *pufM* 系统发育亚群中有 11 个发生过水平基因转移 (HGT)。HGT 不仅发生在同一个纲的物种之间，而且在不同门或亚门的种之间也有发生。不产氧光合细菌的来源环境特征与其系统发育关系密切相关，所有来自有氧栖息地的种和来自缺氧栖息地的种分别聚类在不同的进化枝上。古老的不产氧光合细菌种之间的基因水平转移以及随后它们对各自生态位长期的适应性进化可能导致了不

产氧光合细菌与环境间协同进化的现象。

其次，我们在上述基于不可培养的分子生物学方法研究浮游细菌多样性的基础上，进一步通过培养的方法分析了含色素的有色异养细菌（PHB）这类有重要生态意义的海洋浮游细菌类群的多样性、丰度和光吸收特性。在中国海的近岸、大陆架以及邻近太平洋进行的纯系分离和初步鉴定的研究表明，PHB 的丰度和占总可培养细菌数的比例从近岸到外海逐渐降低，在长江口的值最高，分别为 9.9×10^3 个细胞/毫升和 39.6%。在垂直梯度上，PHB 仅分布于真光层，丰度和比重的最高值均出现在表层。对分离得到的 247 株有色细菌进行 RFLP 筛选和 16S rRNA 基因测序的结果表明，它们覆盖了 6 个细菌类群： α -变形细菌、 γ -变形细菌、*Actinobacteria*、*Bacilli*、*Flavobacteria* 和 *Sphingobacteria*，囊括了 25 个属。不同海区的优势类群和种属分布各有差异；同时，不同种属的菌株颜色也各异，包括金黄色、黄色、红色、粉红和橘色等。细胞色素吸收光的波长介于 450~550 nm 之间。这些结果表明 PHB 广泛分布于海洋环境中，并有很高的多样性，它们在海洋生态系统中对光的吸收和利用有着独特而重要的作用。

最后，对海洋微型生物多样性研究中的一些关键性的方法学问题进行了探讨，并建立了两种新的分析方法或标准，包括：通过对数据库中不产氧光合细菌纯系来源的 *pufM* 序列进行以距离为基础的聚类，并将聚类关系与序列来源物种的分类地位相对应，建立了一套种、属、门（或亚门）上的 *pufM* 序列距离标准，分别为 0.06、0.15 和 0.48；通过结合脉冲场凝胶电泳（PFGE）和 PCR 扩增的技术，建立了一种新的准确估算细菌基因组中 16S rRNA 基因拷贝数的方法。

关键词：海洋浮游细菌；*Cytophaga-Flavobacteria*；浮游古菌；固碳变形细菌；好氧不产氧光合异养细菌（AAPB）；有色异养细菌；16S rRNA 基因；*pufM* 基因；*rbcL* 基因；系统发育分析；遗传多样性；中国海；大洋；弱光层；环境适应机制；协同进化；PFGE

Microbial diversity and Environmental Adaptation Mechanisms in Typical Marine Environments

Abstract

Planktonic bacteria are the most abundant microbes in marine ecosystems. They have extremely high genetic and metabolic diversity and thus play a significant role in marine biogeochemical cycles. Here we attempted to shed new light on the diversity, environmental adaptation mechanisms, and co-evolution with environments of some typical bacterioplankton groups in China seas and global oceans. These bacterial groups play a special role in marine carbon cycling and light utilization but received little attention, including: *Cytophaga-Flavobacteria* (CF), which are proficient in degrading high molecular weight particulate organic carbon; planktonic *Archaea*, the metabolic potential and ecological functions of which are largely unknown; planktonic *Proteobacteria* with the ability of fixing CO₂; and mixotrophic aerobic anoxygenic photosynthetic bacteria (AAPB) and pigmented heterotrophic bacteria (PHB), which can facultatively utilize or absorb light.

Firstly, we chose the Yangtze River estuary (YRE) in the East China Sea (ECS) as a typical eutrophic water and systematically investigated the diversity pattern of CF, planktonic archaea, and CO₂ fixation proteobacteria there. CF cluster is one of the most abundant bacterial groups in the ocean. A newly designed CF specific 16S rRNA gene primer pair was used to construct two clone libraries from YRE and ECS bacterioplankton samples. An abundant CF diversity and a distinct diversity distribution pattern were revealed. Seventy partial 16S rRNA gene sequences of CF were clustered into 26 subgroups, including 7 subgroups that distributed only in the ECS, 17 subgroups that distributed only in the YRE, and 2 subgroups that were shared by both stations. In comparison, the CF species diversity was high in the ECS, but the CF subgroup diversity was high in the YRE. Cosmopolitan subgroups dominated both stations. Subgroups with the soil, coast, and freshwater origins were detected only in the YRE, in accordance with the geographical location and environmental features there. The adaptation of CF species to the two completely different stations with respect to hydrology and the development and succession of local CF populations resulted in these significant differences between the YRE and ECS CF communities.

By the construction of two archaeal 16S rRNA gene clone libraries from YRE bacterioplankton samples, we also revealed the presence of planktonic archaea in typical Chinese marine waters, which has been found to be cosmopolitan in marine ecosystems. Sequencing of the clones in 21 unique RFLP patterns showed the two archaeal groups: Marine Group I (MG I) that was affiliated with *Crenarchaeota* and Marine Group II (MG II) that was affiliated with *Euryarchaeota*. MG I dominated both libraries. A large part of sequences were most related to uncultured archaea, including 2 sequences that were in 98% similarity with a nitrification marine archaeon *Nitrosopumilus maritimus*, suggesting that archaea may play an important role in estuarine ecosystems.

Proteobacteria with the potential of fixing CO₂ is another understudied group in bacterioplankton. Here we for the first time successfully designed two primer pairs targeting the Form I and Form II *rbcL* genes (encoding RubisCO large subunit), respectively, and applied them to the inshore and offshore samples from the ECS. Our results showed that the Form I *rbcL* diversity was high at the inshore station with high salinity and low productivity, while the Form II *rbcL* diversity was high at the offshore station with low salinity and high productivity. A part of sequences showed a low similarity (60~78%) with their nearest neighbors in the GenBank database. The results indicate the presence of carbon-fixing proteobacteria in the ECS and suggest that we should draw more attention to their potential contribution in marine carbon cycling.

Secondly, we focused on a typical functional bacterial group in marine bacterioplankton – AAPB and systematically studied their diversity pattern in global oceans and co-evolution with environments. The diversity data of an AAPB marker gene, *pufM*, was collected from the surface seawater of the Pacific, Atlantic, and Indian oceans, and Chinese marginal seas as well. The results showed that AAPB populations were highly diverse in global oceans and, more importantly, their diversity decreased with the increased Chl.*a* concentrations. This finding indicated that AAPB diversity and abundance followed a reverse trend from oligotrophic to eutrophic oceans.

Depth profile of AAPB diversity was also investigated in three stations in the surface and upper twilight zones of the central Pacific, Atlantic, and Indian oceans by constructing six *pufM* gene clone libraries. The high AAPB diversity in oligotrophic oceans was further confirmed. Furthermore, we for the first time obtained abundant *pufM* sequences (136 sequences within 37 OTU) from deep twilight zones (200 m). Phylogenetic analysis showed that deep AAPB populations were diverse too, covering

all the subgroups that were found in surface stations. They had a lower diversity, GC and GC₃ contents but a higher dominance than surface stations. These results support the hypothesis that AAPB can utilize dim light to acquire energy and thus distribute into the layers below euphotic zones.

To further probe the co-evolutionary relationship between anoxygenic photosynthetic bacteria and environment, we retrieved the *pufM* sequences and related source information of 89 anoxygenic photosynthetic bacterial cultures from public database. Phylogenetic analysis revealed that horizontal gene transfer (HGT) occurred in 11 of total 21 *pufM* phylogenetic subgroups. HGT occurred among species within not only the same class but also different phyla or subphyla. Source environmental features of anoxygenic photosynthetic bacteria were closely related to their phylogeny. Species from oxic and anoxic environments clustered into separate and distinct clades in the phylogenetic tree. HGT between ancient anoxygenic photosynthetic bacteria and long-term adaptive evolution into different niches may result in such coevolutionary events.

Then, in addition to the analysis of bacterial diversity by culture-independent molecular methods, we also employed culture-dependent methods to study the diversity, abundance, and light absorption features of pigmented heterotrophic bacteria (PHB), which is another interesting bacteria group with respect to light utilization. PHB cultures were isolated from the coast and shelf of China seas and adjacent Pacific and were identified by 16S rRNA gene analysis. The results showed that PHB abundance and their percentage in total cultivable bacterial amount decreased from coast to open ocean with the highest values (9.9×10^3 cells mL⁻¹ and 39.6%, respectively) occurring at the YRE station. In the depth profiles, PHB were only distributed in the euphotic zones and the highest values of both abundance and percentage proportion occurred at the surface stations. Total 247 PHB isolates were screened with 16S rRNA gene PCR-RFLP and then sequenced. The PHB strains we obtained covered 6 major bacterial groups (*α-Proteobacteria*, *γ-Proteobacteria*, *Actinobacteria*, *Bacilli*, *Flavobacteria*, and *Sphingobacteria*) and 25 genera. Dominant group differed among different stations, and different genus showed different colors, including golden yellow, yellow, red, pink, and orange etc. Cellular pigments absorbed the light between 450~550 nm. These results indicated that PHB were widely distributed in marine environments with high diversity. They may play a unique and important role in marine ecosystems due to the absorption

and utilization of light.

Finally, we established a set of new criterion for clustering *pufM* gene sequence by distance methods and a new method for assessing 16S rRNA gene copy number in bacterial genomes by combining PFGE and PCR amplification. The cutoff for discriminating species, genus, and (sub-)phylum based on *pufM* gene sequences were 0.06, 0.15, and 0.48, respectively.

Keywords: Marine Bacterioplankton; *Cytophaga-Flavobacteria*; Planktonic Archaea; Carbon-fixation *Proteobacteria*; Aerobic Anoxygenic Photosynthetic Bacteria (AAPB); Pigmented Heterotrophic Bacteria; 16S rRNA gene; *pufM*; *rbcL*; Phylogenetic analysis; Genetic Diversity; China Seas; Open Oceans; Twilight Zones; Environmental Adatation Mechanisms; Co-evolution; PFGE

目 录

摘要.....	I
---------	---

Abstract	IV
----------------	----

第一部分 综述与论文设计

第一章 前言.....	1
--------------------	----------

1.1 海洋异养细菌丰富的遗传多样性	1
1.1.1 异养细菌	1
1.1.2 好氧不产氧光合异养细菌 (AAPB)	4
1.1.3 含视紫质 (Proteorhodopsin, PR) 变形细菌	6
1.1.4 浮游古菌 (Planktonic Archaea)	7
1.2 海洋微生物多样性研究近二十年的主要进展	9
1.2.1 浮游细菌	9
1.2.2 AAPB	14
1.2.3 含视紫质细菌	15
1.2.4 浮游古菌	16
1.2.5 环境基因组学	19
1.2.6 新的开拓性研究方法	20
1.3 本论文的设计与研究意义	22

第二部分 典型浮游细菌类群在中国典型海域的多样性

第二章 浮游 <i>Cytophaga-Flavobacteria</i> 类群在东海的多样性	29
--	-----------

2.1 研究背景	30
2.2 材料和方法	31
2.2.1 采样和环境参数的测定	31
2.2.2 16S rRNA 基因克隆文库的构建	32
2.2.3 测序和系统发育分析	34
2.3 结果	34
2.3.1 东海 CF 类群 16S rRNA 基因序列的总体分析	34
2.3.2 CF 亚群在长江口和东海的多样性	35
2.4 讨论	38
2.4.1 长江口与东海 CF 的多样性分布模式	38
2.4.2 CF 亚群的来源环境分析	40
2.5 小结	42

第三章 浮游古菌在长江口的多样性	43
-------------------------------	-----------

3.1 研究背景	44
3.2 材料和方法	44
3.2.1 采样和环境参数的测定	45
3.2.2 DNA 提取和克隆文库构建	45

3.2.3 选择用于 PCR-RFLP 分析的内切酶	45
3.2.4 克隆文库的 PCR-RFLP 筛选和数据分析	47
3.2.5 测序与系统发育分析	47
3.3 结果	48
3.3.1 采样站位的环境参数	48
3.3.2 PCR-RFLP 筛选克隆文库的限制性内切酶的确定	48
3.3.3 克隆文库的 PCR-RFLP 筛选和带型的统计分析	49
3.3.4 古菌 16S rRNA 基因序列的系统发育分析	53
3.3.5 古菌 16S rRNA 基因序列的同源性分析	53
3.4 讨论	54
第四章 具有固碳能力的浮游变形细菌在东海的多样性	57
4.1 研究背景	58
4.2 材料和方法	59
4.2.1 样品采样和站位环境参数	59
4.2.2 引物设计	60
4.2.3 克隆文库构建	61
4.2.4 DGGE 筛选克隆子	61
4.2.5 测序与系统发育分析	62
4.3 结果	62
4.3.1 引物设计和最佳 PCR 扩增条件的确立	62
4.3.2 I 型和 II 型 <i>rbcL</i> 基因克隆文库的构建和分析	63
4.3.3 <i>rbcL</i> 氨基酸序列的系统发育分析	63
4.3.4 <i>rbcL</i> 氨基酸序列的保守位点分析	67
4.4 讨论	68
4.4.1 引物设计和克隆文库的筛选方法	68
4.4.2 多样性分析及其与环境因子之间的关系	68

第三部分 典型功能细菌类群 AAPB 在全球海洋的多样性

第五章 全球表层海洋 AAPB 多样性的分布模式	70
5.1 研究背景	71
5.2 材料和方法	71
5.3 结果与讨论	73
5.4 小结	77
第六章 AAPB 在大洋表层和弱光层上层水体中的多样性	78
6.1 研究背景	79
6.2 材料和方法	80
6.2.1 采样站位和环境参数	80
6.2.2 样品采集和群落 DNA 提取	80
6.2.3 AAPB 多样性分析的目标基因和引物	82
6.2.4 PCR 扩增目标基因片段以及克隆文库分析	82
6.2.5 测序与系统发育分析	82
6.2.6 统计分析	83
6.3 结果	84

6.3.1 采样情况与环境参数	84
6.3.2 <i>pufM</i> 克隆文库分析	84
6.3.3 序列统计分析	85
6.3.4 AAPB 多样性模式	87
6.4 讨论	88
6.4.1 AAPB 多样性的研究方法	91
6.4.2 全球海洋 AAPB 生物地理分布的新认识	94
6.4.3 弱光层 AAPB 的适应和进化启示	97
6.5 小结	98
第七章 不产氧光合细菌与环境的协同进化关系研究	100
7.1 研究背景	101
7.2 材料和方法	102
7.2.1 收集 APB 菌株的 <i>pufM</i> 序列和来源环境信息	102
7.2.2 <i>pufM</i> 序列的系统发育分析	105
7.3 结果与讨论	105
7.3.1 <i>pufM</i> 系统发育树的构建与分析	105
7.3.2 <i>pufM</i> 系统发育和来源环境之间的关系分析	108
第四部分 海洋可培养的有色异养细菌的多样性	
第八章 中国海及邻近大洋海域可培养有色异养细菌的多样性分布和生理特征 ..	111
8.1 研究背景	112
8.2 材料和方法	113
8.2.1 采样站位	113
8.2.2 总异养细菌计数	113
8.2.3 CFU 和 PHB 计数	113
8.2.4 16S rRNA 基因的测序与系统发育分析	113
8.2.5 丙酮-甲醇提取物的色素分析	115
8.3 结果	115
8.3.1 CFU 和 PHB 的空间分布	115
8.3.2 总细菌数、CFU 和 PHB 的垂直剖面	116
8.3.3 16S rRNA 基因序列分析	118
8.3.4 色素吸收光谱分析	124
8.4 讨论	124
8.4.1 海洋中 PHB 的丰度分布	124
8.4.2 PHB 的遗传多样性	125
第五部分 方法学研究	
第九章 基于 <i>pufM</i> 序列距离的聚类方法评估不产氧光合细菌的多样性	129
9.1 研究背景	130
9.2 材料和方法	130
9.2.1 <i>pufM</i> 序列的收集	130
9.2.2 <i>pufM</i> 序列的距离聚类方法	131

9.3 结果	132
9.3.1 纯培养 <i>pufM</i> 序列分析及 cutoff 值的确定	132
9.3.2 直接来源于环境的 <i>pufM</i> 序列的分析	137
9.3.3 <i>pufM</i> 序列在 <i>Culture</i> 和 <i>EnvClone</i> 两种类群间的距离	137
9.4 讨论	137
9.4.1 评估基于序列距离的聚类方法和 cutoff 值	137
9.4.2 自然界中 APB 的多样性	140
第十章 结合脉冲场电泳与 PCR 扩增的方法估算细菌基因组中 16S rRNA 基因的拷贝数	142
10.1 研究背景	143
10.2 材料和方法	143
10.3 结果和讨论	144
10.4 小结	146
第六部分 本论文的主要结论和创新点	148
参考文献	150
附录 1 缩写列表	172
附录 2 本论文的技术路线示意图	173
附录 3 常用分子生物学软件、程序及使用方法	174
附录 4 博士期间参予课题和论文发表情况	183

CONTENTS

Chinese Abstract	I
------------------------	---

Abstract	IV
----------------	----

Part I Review and Thesis Design

Chapter 1 Preface	1
1.1 Abundant diversity of marine heterotrophic bacteria	1
1.1.1 Heterotrophic bacteria	1
1.1.2 Aerobic anoxygenic photosynthetic bacteria	4
1.1.3 Rhodopsin-containing <i>Proteobacteria</i>	6
1.1.4 Planktonic <i>Archaea</i>	7
1.2 Major advances in marine microbial diversity studies during the past twenty years	9
1.2.1 Bacterioplankton	9
1.2.2 AAPB	14
1.2.3 Proteorhodopsin (PR) bacteria	15
1.2.4 Planktonic <i>Archaea</i>	16
1.2.5 Environmental genomics	19
1.2.6 Breakthroughs in the research methods	20
1.3 Design of this thesis and research significance	22

Part II Diversity of typical groups of bacterioplankton in typical areas of China seas

Chapter 2 Diversity of planktonic <i>Cytophaga-Flavobacteria</i> cluster in the East China Sea (ECS)	29
2.1 Background	30
2.2 Materials and Methods	31
2.2.1 Sampling and environmental variables	31
2.2.2 Construction of CF 16S rRNA gene clone libraries	32
2.2.3 Sequencing and phylogenetic analysis	34
2.3 Results	34
2.3.1 Overall analysis of CF 16S rRNA gene sequences	34
2.3.2 Diversity of CF subgroups in the Yangtze River estuary (YRE) and the ECS	35
2.4 Discussion	38
2.4.1 Distribution pattern of CF diversity in the YRE and ECS	38
2.4.2 Analysis of the source features of CF subgroups	40
2.5 Conclusion	42
Chapter 3 Diversity of planktonic <i>Archaea</i> in the YRE	43
3.1 Background	44
3.2 Materials and Methods	44

3.2.1 Sampling and environmental variables	45
3.2.2 DNA extraction and clone library construction	45
3.2.3 Enzymes for PCR-RFLP analysis	45
3.2.4 PCR-RFLP screening of clone libraries and data statistics	47
3.2.5 Sequencing and phylogenetic analysis	47
3.3 Results	48
3.3.1 Analysis of environmental variables	48
3.3.2 Determination of enzymes for PCR-RFLP screening	48
3.3.3 PCR-RFLP screening and statistical analysis of RFLP bands	49
3.3.4 Phylogenetic analysis of archaeal 16S rRNA gene sequences	53
3.3.5 Similarity analysis of archaeal 16S rRNA gene sequences	53
3.4 Discussion	54
Chapter 4 Diversity of the planktonic proteobacteria with the ability of fixing carbon in the ECS	57
4.1 Background	58
4.2 Materials and Methods	59
4.2.1 Sampling and environmental variables	59
4.2.2 Primer design	60
4.2.3 Construction of clone library	61
4.2.4 DGGE screening of clones	61
4.2.5 Sequencing and phylogenetic analysis	62
4.3 Results	62
4.3.1 Primer design and determination of optimal PCR amplification conditions	62
4.3.2 Clone library analysis of Form I and II <i>rbcL</i> genes	63
4.3.3 Phylogenetic analysis of the amino acid sequences of <i>rbcL</i> genes	63
4.3.4 Analysis of conserved sites in the amino acid sequences of <i>rbcL</i> genes	67
4.4 Discussion	68
4.4.1 Assessment of primers and screening methods for clone libraries	68
4.4.2 Diversity and its correlation with environmental variables	68
Part III Diversity of a typical functional bacterial group – AAPB in global oceans	
Chapter 5 Diversity of AAPB in global surface oceans	70
5.1 Background	71
5.2 Materials and Methods	71
5.3 Results and Discussion	73
5.4 Conclusion	77
Chapter 6 Diversity of AAPB in surface oceans and upper twilight zones	78
6.1 Background	79
6.2 Materials and Methods	80
6.2.1 Sampling stations and environmental variables	80
6.2.2 Sampling and DNA extraction	80
6.2.3 Target gene and primer for AAPB diversity analysis	82
6.2.4 PCR and clone library analysis	82
6.2.5 Sequencing and phylogenetic analysis	82
6.2.6 Statistical analysis	83

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库